

St Joseph's Catholic Primary School Anaphylaxis Procedures.

Adopted by St. Joseph's Catholic Primary School on 27th September 2023. Procedures to be read in conjunction with 'Managing Medicines in School Policy'. These procedures will be reviewed annually.

1. Introduction

Anaphylaxis is a serious, life-threatening allergic reaction. It is at the extreme end of the allergic spectrum. The whole body is affected often within minutes of exposure to the allergen, but sometimes it can be hours later. Causes can include foods, insect stings, and drugs.

Most healthcare professionals consider an allergic reaction to be anaphylaxis when it involves difficulty breathing or affects the heart rhythm or blood pressure. Anaphylaxis symptoms are often referred to as the ABC symptoms (Airway, Breathing, Circulation).

It is possible to be allergic to anything which contains a protein, however most people will react to a fairly small group of potent allergens.

2. Role and responsibilities

Parent Responsibilities

- On entry to the school, it is the parent's responsibility to inform Senior Administrator of any allergies. This information should include all previous serious allergic reactions, history of anaphylaxis and details of all prescribed medication.
- Parents are to supply a copy of their child's Allergy Action Plan
- Parents are responsible for ensuring any required medication is supplied, in date and replaced as necessary.
- Parents are requested to keep the school up to date with any changes in allergy management. The Allergy Action Plan will be kept updated accordingly.

Staff Responsibilities

- All staff will complete anaphylaxis training. Training is provided for all staff on a yearly basis and on an ad-hoc basis for any new members of staff.
- Staff must be aware of the pupils in their care (regular or cover classes) who have known allergies as an allergic reaction could occur at any time and not just at mealtimes. Any food-related activities must be supervised with due caution.
- Staff leading school trips will ensure they carry all relevant emergency supplies. Trip leaders will check that all pupils with medical conditions, including allergies, carry their medication. Pupils unable to produce their required medication will not be able to attend the excursion.
- Senior Administrator will ensure that the up-to-date Allergy Action Plan is kept with the pupil's medication.
- It is the parent's responsibility to ensure all medication in date however the Senior Administrator will check medication kept at school on a termly basis and send a reminder to parents if medication is approaching expiry.

- Senior Administrator keeps a register of pupils who have been prescribed an adrenaline auto-injector (AAI) and a record of use of any AAI(s) and emergency treatment given.

Pupil Responsibilities

- Pupils are encouraged to have a good awareness of their symptoms and to let an adult know as soon as they suspect they are having an allergic reaction.

3. Allergy action plans

Allergy action plans are designed to function as individual healthcare plans for children with food allergies, providing medical and parental consent for schools to administer medicines in the event of an allergic reaction, including consent to administer a spare adrenaline auto-injector.

It is the parent/carer's responsibility to complete the allergy action plan with help from a healthcare professional and provide this to the school.

4. Emergency Treatment and Management of Anaphylaxis

What to look for:

Symptoms usually come on quickly, within minutes of exposure to the allergen.

Mild to moderate allergic reaction symptoms may include:

- a red raised rash (known as hives or urticaria) anywhere on the body
- a tingling or itchy feeling in the mouth
- swelling of lips, face or eyes
- stomach pain or vomiting.

More serious symptoms are often referred to as the ABC symptoms and can include:

- AIRWAY - swelling in the throat, tongue or upper airways (tightening of the throat, hoarse voice, difficulty swallowing).
- BREATHING - sudden onset wheezing, breathing difficulty, noisy breathing.
- CIRCULATION - dizziness, feeling faint, sudden sleepiness, tiredness, confusion, pale clammy skin, loss of consciousness.

The term for this more serious reaction is anaphylaxis. In extreme cases there could be a dramatic fall in blood pressure. The person may become weak and floppy and may have a sense of something terrible happening. This may lead to collapse and unconsciousness and, on rare occasions, can be fatal.

If the pupil has been exposed to something they are known to be allergic to, then it is more likely to be an anaphylactic reaction. Anaphylaxis can develop very rapidly, so a treatment is needed that works rapidly. **Adrenaline** is the mainstay of treatment, and it starts to work within seconds.

What does adrenaline do?

- It opens up the airways
- It stops swelling

- It raises the blood pressure

As soon as anaphylaxis is suspected, adrenaline must be administered without delay.

Action:

- Keep the child where they are, call for help and do not leave them unattended.
- **LIE CHILD FLAT WITH LEGS RAISED** – they can be propped up if struggling to breathe but this should be for as short a time as possible.
- **USE ADRENALINE AUTO-INJECTOR WITHOUT DELAY** and note the time given. AAI should be given into the muscle in the outer thigh. Specific instructions vary by brand – always follow the instructions on the device.
- CALL **999** and state **ANAPHYLAXIS (ana-fil-axis)**.
- If no improvement after 5 minutes, administer second AAI.
- If no signs of life commence CPR.
- Call parent/carer as soon as possible.

Whilst you are waiting for the ambulance, keep the child where they are. Do not stand them up, or sit them in a chair, even if they are feeling better. This could lower their blood pressure drastically, causing their heart to stop. All pupils must go to hospital for observation after anaphylaxis even if they appear to have recovered as a reaction can reoccur after treatment.

5. Supply, storage and care of medication

A child's anaphylaxis kit is kept safely in the classroom, and accessible to all staff.

Medication should be stored in a suitable container and clearly labelled with the pupil's name. The pupil's medication storage container should contain:

- Two AAIs i.e. EpiPen® or Jext® or Emerade®
- An up-to-date allergy action plan
- Antihistamine as tablets or syrup (if included on allergy action plan)
- Spoon if required
- Asthma inhaler (if included on allergy action plan).

It is the responsibility of the child's parents to ensure that the anaphylaxis kit is up-to-date and clearly labelled, however the Senior Administrator will check medication kept at school on a termly basis and send a reminder to parents if medication is approaching expiry.

AAIs should be stored at room temperature, protected from direct sunlight and temperature extremes. AAIs are single use only and must be disposed of as sharps. Used AAIs can be given to ambulance paramedics on arrival or can be disposed of in a pre-ordered sharps bin.

The Senior Administrator is responsible for checking if the spare medication is in date monthly and to replace as needed. Written parental permission for use of the spare AAIs is included in the pupil's allergy action plan.

If anaphylaxis is suspected **in an undiagnosed individual** call the emergency services and state, you suspect ANAPHYLAXIS. Follow advice from them as to whether administration of the spare AAI is appropriate.

6. Staff Training

Senior Administrator is responsible for co-ordinating staff anaphylaxis training and the upkeep of the school's anaphylaxis policy.

All staff will complete online AllergyWise anaphylaxis training at the start of every new academic year (supplied by NHS nurse team). Training is also available on an ad-hoc basis for any new members of staff. Staff working directly with identified children will have a six-monthly refresher using webinar from Anaphylaxis UK.

Training includes:

- Regular support and guidance from 'Kelly Lambert' DCC, Senior Advisor Physical and Medical Needs service is provided when required.
- Knowing the common allergens and triggers of allergy
- Spotting the signs and symptoms of an allergic reaction and anaphylaxis. Early recognition of symptoms is key, including knowing when to call for emergency services
- Administering emergency treatment (including AAIs) in the event of anaphylaxis – knowing how and when to administer the medication/device
- Measures to reduce the risk of a child having an allergic reaction e.g. allergen avoidance, knowing who is responsible for what
- Managing allergy action plans and ensuring these are up to date
- A practical session using trainer devices (these can be obtained from the manufacturers' websites: www.epipen.co.uk and www.jext.co.uk and www.emerade-bausch.co.uk)
- Whole school assembly is delivered each year to children. Materials provided by Anyaphlaxis UK.

7. Catering

All food businesses (including school caterers) must follow the Food Information Regulations 2014 which states that allergen information relating to the 'Top 14' allergens must be available for all food products.

The school menu is available for parents to view in advance with all ingredients listed and allergens highlighted on the school website at <https://www.sjcpschool.co.uk/parents/school-meals>

The Senior Administrator will inform the Lunchtime Manager of pupils with food allergies. A photograph of children with allergies is displayed in the kitchen servery. School's Senior Administrator ensures this is reviewed termly or when a new child starts school who has allergies.

The school adheres to the following Department of Health guidance recommendations:

- Bottles, other drinks and lunch boxes provided by parents for pupils with food allergies should be clearly labelled with the name of the child for whom they are intended.
- If food is purchased from the school canteen/tuck shop, parents should check the appropriateness of foods by speaking directly to the catering manager.
- Where food is provided by the school, staff should be educated about how to read labels for food allergens and instructed about measures to prevent cross contamination during the handling, preparation and serving of food. Examples include preparing food for children with food allergies first; careful cleaning (using warm soapy water) of food preparation areas and utensils. For further information, parents/carers are encouraged to liaise with the Catering Manager.
- Food should not be given to primary school age food-allergic children without parental engagement and permission (e.g. birthday parties, food treats).
- Use of food in crafts, cooking classes, science experiments and special events (e.g. fetes, assemblies, cultural events) needs to be considered and may need to be restricted/risk assessed depending on the allergies of particular children and their age.

8. School trips

Staff leading school trips will ensure they carry all relevant emergency supplies. Trip leaders will check that all pupils with medical conditions, including allergies, carry their medication. Pupils unable to produce their required medication will not be able to attend the excursion.

All the activities on the school trip will be risk assessed to see if they pose a threat to allergic pupils and alternative activities planned to ensure inclusion.

Overnight school trips should be possible with careful planning and a meeting for parents with the lead member of staff planning the trip should be arranged. Staff at the venue for an overnight school trip should be briefed early on that an allergic child is attending and will need appropriate food (if provided by the venue).

Sporting Excursions

The school will ensure that the P.E. teacher/s are fully aware of the situation. The school being visited will be notified that a member of the team has an allergy when arranging the fixture. A member of staff trained in administering adrenaline will accompany the team. If another school feels that they are not equipped to cater for any food-allergic child, the school will arrange for the child to take alternative/their own food.

9. Allergy awareness and nut bans

School supports the approach advocated by Anaphylaxis UK towards nut bans/nut free schools. They would not necessarily support a blanket ban on any particular allergen in any establishment, including in schools. This is because nuts are only one of many allergens that could affect pupils, and no school could guarantee a truly allergen free environment for a child living with food allergy. They advocate instead for schools to adopt a culture of allergy awareness and education.

A 'whole school awareness of allergies' is a much better approach, as it ensures teachers, pupils and all other staff are aware of what allergies are, the importance of avoiding the pupils' allergens, the signs & symptoms, how to deal with allergic reactions and to ensure policies and procedures are in place to minimise risk.

10. Risk Assessment

School will conduct a detailed individual risk assessment for all new joining pupils with allergies and any pupils newly diagnosed, to help identify any gaps in our systems and processes for keeping allergic children safe.

11. Useful Links

Anaphylaxis UK Safer Schools Programme -

<https://www.anaphylaxis.org.uk/education/safer-schools-programme/>

AllergyWise for Schools (including certificate) online training -

<https://www.allergywise.org.uk/p/allergywise-for-schools1>

BSACI Allergy Action Plans - <https://www.bsaci.org/professional-resources/resources/paediatric-allergy-action-plans/>

Spare Pens in Schools - <http://www.sparepensinschools.uk>

Department for Education Supporting pupils at school with medical conditions -

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/803956/supporting-pupils-at-school-with-medical-conditions.pdf

Department of Health Guidance on the use of adrenaline auto-injectors in schools -

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/645476/Adrenaline_auto_injectors_in_schools.pdf

Food allergy quality standards (The National Institute for Health and Care Excellence, March 2016) <https://www.nice.org.uk/guidance/qs118>

Anaphylaxis: assessment and referral after emergency treatment (The National Institute for Health and Care Excellence, 2020)

<https://www.nice.org.uk/guidance/cg134?unlid=22904150420167115834>

September 2024